

# *Latest on the Proton Charge Radius from the PRad Experiment*

*Haiyan Gao*

*Duke University and Duke Kunshan  
University*




# Lepton scattering: powerful microscope!



- Clean probe of hadron structure
- Electron (lepton) vertex is well-known from QED
- One-photon exchange dominates, *higher-order exchange diagrams are suppressed (two-photon physics)*
- *Vary the wave-length of the probe to view deeper inside*


$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4E^2 \sin^4 \frac{\theta}{2}} \frac{E'}{E} \left( \frac{G_E^2 + \tau G_M^2}{1 + \tau} \cos^2 \frac{\theta}{2} + 2\tau G_M^2 \sin^2 \frac{\theta}{2} \right) \quad \tau = -q^2 / 4M^2$$



Virtual photon 4-momentum

$$q = k - k' = (\vec{q}, \omega)$$

$$Q^2 = -q^2$$



# What is inside the proton/neutron?

1933: Proton's magnetic moment



Nobel Prize  
In Physics 1943

Otto Stern

"for ... and for his discovery of the magnetic moment of the proton".

$$g \neq 2$$

1960: Elastic e-p scattering




Nobel Prize  
In Physics 1961



Robert Hofstadter

"for ... and for his thereby achieved discoveries concerning the structure of the nucleons"

Form factors → Charge distributions



1969: Deep inelastic e-p scattering



Nobel Prize in Physics 1990

Jerome I. Friedman, Henry W. Kendall, Richard E. Taylor

"for their pioneering investigations concerning deep inelastic scattering of electrons on protons ...".

Jian-Wei Qiu

1974: QCD Asymptotic Freedom



Nobel Prize in Physics 2004

David J. Gross, H. David Politzer, Frank Wilczek


"for the discovery of asymptotic freedom in the theory of the strong interaction".

# *Proton Charge Radius*

- An important property of the nucleon
  - Important for understanding how QCD works
  - Challenge to Lattice QCD (exciting new results, Alexandrou et al.)
  - An important physics input to the bound state QED calculations, affects muonic H Lamb shift ( $2S_{1/2} - 2P_{1/2}$ ) by as much as 2%
- Electron-proton elastic scattering to determine electric form factor (Nuclear Physics)


$$\sqrt{\langle r^2 \rangle} = \sqrt{-6 \frac{dG(q^2)}{dq^2} \Big|_{q^2=0}}$$

- Spectroscopy (Atomic physics)
  - Hydrogen Lamb shift
  - Muonic Hydrogen Lamb shift





# Proton Charge Radius Puzzle



Electron:  $0.8751 \pm 0.0061$  fm

Muon:  $0.8409 \pm 0.0004$  fm

- p Lamb shift measurements by CREMA (2010, 2013)
  - Unprecedented precision,  $<0.1\%$

# Unpolarized electron-nucleon scattering

## (Rosenbluth Separation)

- Elastic e-p cross section



$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2 \cos^2 \frac{\theta}{2}}{4E^2 \sin^4 \frac{\theta}{2}} \frac{E'}{E} \left( \frac{G_E^p{}^2 + \tau G_M^p{}^2}{1 + \tau} + 2\tau G_M^p{}^2 \tan^2 \frac{\theta}{2} \right)$$

$$= \sigma_M f_{rec}^{-1} \left( A + B \tan^2 \frac{\theta}{2} \right)$$

- At fixed  $Q^2$ , fit  $d\sigma/d\Omega$  vs.  $\tan^2(\theta/2)$ 
  - Measurement of absolute cross section
  - Dominated by either  $G_E$  or  $G_M$**

- Low  $Q^2$  by  $G_E$
- High  $Q^2$  by  $G_M$

$G_E$  or  $G_M$



$$\sigma_R = \tau G_M^2 + \epsilon G_E^2$$

$$\tau = \frac{Q^2}{4M^2}$$


$$\epsilon = (1 + 2(1 + \tau) \tan^2 \frac{\theta}{2})^{-1}$$

# Electron-proton elastic scattering with longitudinally polarized electron beam and recoil proton polarization measurement

## Polarization Transfer



$$\frac{G_E^p}{G_M^p}$$





- Recoil proton polarization

$$\frac{G_E^p}{G_M^p} = -\frac{P_t E + E'}{P_l 2M} \tan \frac{\theta}{2}$$

- Focal Plane Polarimeter

- recoil proton scatters off secondary  $^{12}\text{C}$  target
- $P_t$ ,  $P_l$  measured from  $\varphi$  distribution
- $P_b$ , and analyzing power cancel out in ratio





Focal-plane polarimeter

# Asymmetry Super-ratio Method

## Polarized electron-polarized proton elastic scattering


- Polarized beam-target asymmetry

$$A_{exp} = P_b P_t \frac{-2\tau v_{T'} \cos \theta^* G_M^p{}^2 + 2\sqrt{2\tau(1+\tau)} v_{TL'} \sin \theta^* \cos \phi^* G_M^p G_E^p}{(1+\tau) v_L G_E^p{}^2 + 2\tau v_T G_M^p{}^2}$$




- Super-ratio

$$R_A = \frac{A_1}{A_2} = \frac{a_1 - b_1 \cdot G_E^p / G_M^p}{a_2 - b_2 \cdot G_E^p / G_M^p}$$



BLAST pioneered the technique, later also used in Jlab Hall A experiment

# Hydrogen Spectroscopy



The absolute frequency of H energy levels has been measured with an accuracy of **1.4 part in  $10^{14}$**  via comparison with an **atomic cesium fountain clock** as a primary frequency standard.

Yields  $R_\infty$  (the most precisely known constant)

Comparing measurements to QED calculations that include corrections for the finite size of the proton provide an **indirect** but very precise value of the **rms proton charge radius**




Proton charge radius effect on the muonic hydrogen Lamb shift is 2%



# Muonic hydrogen Lamb shift at PSI (2010, 2013)




*Nature* **466**, 213-216 (8 July 2010)



2010: new value is  $r_p = 0.84184(67)$  fm

# New PSI results reported in Science 2013



2013:  $r_p = 0.84087(39)$  fm, A. Antognini *et al.*, Science 339, 417 (2013)

# Recent $ep$ Scattering Experiments


Three spectrometer facility of the A1 collaboration:



- Large amount of overlapping data sets
  - Statistical error  $\leq 0.2\%$
  - Luminosity monitoring with spectrometer
  - $Q^2 = 0.004 - 1.0 \text{ (GeV/c)}^2$
- result:  $r_p = 0.879(5)_{\text{stat}}(4)_{\text{sys}}(2)_{\text{mod}}(4)_{\text{group}}$

J. Bernauer, PRL 105,242001, 2010

## Measurements @ Mainz




5-7 $\sigma$  higher than muonic hydrogen result !

(J. Bernauer)

# JLab Recoil Proton Polarization Experimental

LHRS


- $\Delta p/p_0: \pm 4.5\%$  ,
- out-of-plane:  $\pm 60$  mrad
- in-plane:  $\pm 30$  mrad
- $\Delta\Omega: 6.7$ msr
- QQDQ
- Dipole bending angle  $45^\circ$
- **VDC+FPP**
- $P_p: 0.55 \sim 0.93$  GeV/c



$E_e: 1.192$ GeV  
 $P_b: \sim 83\%$


BigBite

- Non-focusing Dipole
- Big acceptance.
  - $\Delta p: 200-900$ MeV
  - $\Delta\Omega: 96$ msr
- PS + Scint. + **SH**



*X. Zhan et al. Phys. Lett. B 705 (2011) 59-64*  
*C. Crawford et al. PRL98, 052301 (2007)*

# *Proton Charge Radius from recent experiments and analyses*





# Revisits QED Calculations....

An additional 0.31 meV to match CODATA value

Contribution	Value [meV]	Uncertainty [ $10^{-4}$ meV]
Uehling	205.0282	
Källen-Sabry	1.5081	
VP iteration	0.151	
Mixed $\mu - e$ VP	0.00007	
Hadronic VP [21, 23]	0.011	20
Sixth order VP [24]	0.00761	
Whichmann-Kroll	-0.00103	
Virtual Delbrück	0.00135	
Light-by-light	-	10
Muon self-energy and muonic VP (2 <sup>nd</sup> order)	-0.66788	
Fourth order electron loops	-0.00169	
VP insertion in self energy [17]	-0.0055	10
Proton self-energy [18]	-0.0099	
Recoil [17, 43]	0.0575	
Recoil correction to VP (one-photon)	-0.0041	
Recoil (two-photon) [19]	-0.04497	
Recoil higher order [19]	-0.0096	
Recoil finite size [32]	0.013	10
Finite size of order $(Z\alpha)^4$ [32]	$-5.1975(1) r_p^2$	(620)
Finite size of order $(Z\alpha)^5$	$0.0347(30) r_p^3$	(20)
Finite size of order $(Z\alpha)^6$	-0.0005	
Correction to VP	$-0.0109 r_p^2$	
Additional size for VP [19]	$-0.0164 r_p^3$	
Proton polarizability [18, 33]	0.015	40
Fine structure $\Delta E(2P_{3/2} - 2P_{1/2})$	8.352	10
$2P_{3/2}^{F=2}$ hyperfine splitting	1.2724	
$2S_{1/2}^{F=1}$ hyperfine splitting [42], $(-22.8148/4)$	-5.7037	20

Evaluation by Jentschura, Annals Phys. 326, 500 (2011)  
Recent summary by A. Antognini et al., arXiv:1208.2637

Birse and McGovern, arXiv:1206.3030  
0.015(4) meV (proton polarizability)

J.M. Alarcon, et al. 1312.1219  
0.008 meV

G.A. Miller, arXiv:1209.4667

New experiments at HIGS and Mainz on proton polarizabilities

# *Revisits of e-p scattering data (just 2015)*

- Re-analysis of existing proton form factor data
  - D. W. Higinbotham, arXiv:1510.01293: two parameter dipole form fit describes the data at both low  $Q^2$  and high  $Q^2$  well, and the result is consistent with PSI value
  - K. Griffioen, C. Carson, S. Maddox, arXiv:1509.06676: re-analysis of Mainz data, focusing on the low  $Q^2$  part with a polynomial form fit.
  - M. Horbatsch and E. A. Hessels, arXiv:1509.05644: re-analysis of Mainz data, simple fits (one-parameter model, dipole model, linear model) for low  $Q^2$  data, and spline extension to high  $Q^2$  data, these fits can all describe data well, but the extracted radius varies from 0.84 ~ 0.89 fm. So current data is not able to resolve the puzzle.
  - J. Arrington, arXiv:1506.00873: re-analysis of world data, found the previous scattering results might underestimate the uncertainty.
  - Distler, Walcher, and Bernauer, arXiv:1511.00479

*All these studies emphasize even more the importance of low  $Q^2$  e-p scattering data*


## *New Physics or what? - Incomplete list*

- **New physics: new particles**, Barger et al., Carlson and Rislw; Liu and Miller,....**New PV muonic force**, Batell et al.; Carlson and Freid;  
**Extra dimension**: Dahia and Lemos; **Quantum gravity at the Fermi scale** R. Onofrio;.....
- **Contributions to the muonic H Lamb shift**: Carlson and Vanderhaeghen,; Jentschura, Borie, Carroll et al, Hill and Paz, Birse and McGovern, G.A. Miller, J.M. Alarcon, Ji, Peset and Pineda....
- **Higher moments of the charge distribution and Zemach radii**, Distler, Bernauer and Walcher,.....
- J.A. Arrington, G. Lee, J. R. Arrington, R. J. Hill discuss systematics in extraction from ep data, no resolution on discrepancy
- Donnelly, Milner and Hasell discuss interpretation of ep data,.....

**Discrepancy explained by some but others disagree**

- Dispersion relations: Lorentz et al.
- Frame transformation: D. Robson
- **New experiments: Mainz (e-d, ISR), JLab (PRad), PSI (Lamb shift, and MUSE), H Lamb shift**

# The Proton Radius Puzzle (June 2016)



- New, preliminary value for  $r_p$  was reported in PRP-2016 Workshop (Trento, Italy) from ordinary hydrogen
- Consistent with the muonic-hydrogen result !
- Is the Puzzle solved? No, new measurements are needed (spectroscopy, ep-scattering)

# *Update on proton radius puzzle*


- Deuteron radius puzzle
  - Deuteron rms charge radius from muonic deuterium spectroscopy (R. Pohl et al., [Science 353, 6300, 669, 2016](#))
  - $7.5\sigma$  smaller than the CODATA-2010 value, and  $3.5\sigma$  smaller than the value from electronic deuterium spectroscopy (R. Pohl et al., [Metrologia 54, L1, 2017](#))
  - Confirms proton radius puzzle
- Analysis of electron scattering data
  - Focusing on the low-q data yields a consistent result with CREMA's value K. Griffioen, C. Carlson, and S. Maddox. ([Phy. Rev. C 93, 065207, 2016](#))  
D. Higinbotham, A.A. Kabir, V. Lin, D. Meekins, B. Norum, and B. Sawatzky. ([Phys. Rev. C 93, 055207, 2016](#))  
M. Horbatsch and E.A. Hessels. ([Phys. Rev. C 93, 015204, 2016](#))
  - However, I. Sick and D. Trautmann ([Phys. Rev. C 95, 012501\(R\), 2017](#)) claim that the above analyses led to a systematically smaller proton rms-radius because of the ignorance of the correlations from higher moments  $\langle r^{2n} \rangle$



# Deuteron Charge Radius?

- “Proton Charge Radius Puzzle” is still **unsolved** after seven years.
- There is a newly developing “Deuteron Charge Radius Puzzle”

H/D isotope shift:  $r_d^2 - r_p^2 = 3.82007(65) \text{ fm}^2$   
 Muonic deuterium:  $r_d = 2.12562(13)_{\text{exp}}(77)_{\text{theory}} \text{ fm}$   
 Electronic deuterium:  $r_d = 2.14150(450) \text{ fm}$




(R. Pohl, 2017)

- Calls for new independent experiments with possible highest accuracy!
- New ed- cross sections at low  $Q^2$  will be a critical input to reduce theory error in  $r_d$  extracted from  $\mu\text{D}$  spectroscopy.


# Charge Radius of Helium Nuclei

Helium



Electron scattering consistent with  $\mu$ -spectroscopy

# PRad Experimental Setup in Hall B at JLab



- High resolution, large acceptance, hybrid HyCal calorimeter (**PbWO<sub>4</sub>** and **Pb-Glass**)
- Windowless H<sub>2</sub> gas flow target
- Simultaneous detection of elastic and Moller electrons
- Q<sup>2</sup> range of **2x10<sup>-4</sup> – 0.14 GeV<sup>2</sup>**
- XY – veto counters replaced by GEM detector
- Vacuum chamber

Spokespersons: D. Dutta, H. Gao, A. Gasparian, M. Khandaker


Sub 1% measurements:

- (1) ep elastic scattering at Jlab (PRad)
- (2)  $\mu$ p elastic scattering at PSI - 16 U.S. institutions! (MUSE)
- (3) ISR experiments at Mainz


Ongoing H spectroscopy experiments<sup>22</sup>

# PRad Experimental Apparatus

PRad Setup (Side View)




- 8 cm dia x 4 cm long target cell
- 2 mm holes open at front and back kapton foils, allows beam to pass through
- Target thickness:  $\sim 2 \times 10^{18}$  H atoms / cm<sup>2</sup>



# PRad Experimental Apparatus

PRad Setup (Side View)




- 5 m long two stage vacuum chamber, further remove possible background source
- vacuum tank pressure: 0.3 mTorr




# PRad Experimental Apparatus

## PRad Setup (Side View)




- Two large area GEM detectors
- Small overlap region in the middle
- Excellent position resolution ( $72 \mu\text{m}$ )
- Improve position resolution of the setup by  $> 20$  times
- Similar improvement for  $Q^2$  determination at small angle




More details see presentation of X. Bai in session E12

# PRad Experimental Apparatus

## PRad Setup (Side View)





- Hybrid EM calorimeter (HyCal)
  - Inner 1156  $\text{PWO}_4$  modules
  - Outer 576 lead glass modules
- 5.8 m from the target
- Scattering angle coverage:  $\sim 0.6^\circ$  to  $7.5^\circ$
- Full azimuthal angle coverage
- High resolution and efficiency



# HyCal Resolution and Efficiency


- HyCal energy resolution and trigger efficiency extracted using high energy photon beam from Hall B at Jlab
  - > 99.5% trigger efficiency obtained for  $E_\gamma > 500$  MeV, for various parts of HyCal
  - Energy resolution  $\sim 2.5\%$  for  $\text{PbWO}_4$  part, lead glass part about 2.5 time worse




# Performance of GEM Detectors

- GEM detection efficiency measured in both photon beam calibration (**pair production**) and production runs (***ep* and *ee***)
- Using overlap region of GEMs to measure position resolution ( $72 \mu\text{m}$ )


GEM Efficiency in Active Area



Position Resolution




# Preliminary Results:




# Preliminary Elastic $ep$ Cross Section

- Plots show the extracted differential cross section v.s. scattering angle and  $Q^2$ , with 2.2 GeV data in 0.7 ~ 3.5 deg range (very preliminary)
- Statistical error at this stage:  $\sim 0.2\%$  per point
- Systematic errors are conservatively assigned at  $\sim 2\%$  at current stage (shown as shadow area)

$ep$  elastic scattering cross section



$ep$  elastic scattering cross section







# PRad Analysis Status: Event Selection Quality

- Control of background in the PRad experiment.
- Consistency of two practically independent measurements (within the  $\sim 0.2\%$  statistical errors) demonstrates that we **control the background**, and

PRad will reach its goal of sub-percent extraction of the Proton Radius!!!



# *PRad Projected Result with world data*



# *Summary and outlook*

- After several years, the proton charge radius remains puzzling, and perhaps also the deuteron charge radius
- PRad experiment had a successful data taking in May/June 2016
- PRad collaboration is making good progress in data analysis and preliminary cross section results (partial data) announced in June 2017
- Preliminary radius result is anticipated in the fall 2017 –Stay tuned!

**Acknowledgement: the SoLID collaboration, and the PRad Collaboration (supported in part by U.S. Department of Energy under contract number DE-FG02-03ER41231, NSF MRI PHY-1229153)**