



# A test of first order scaling in $N_f=2$ QCD: progress report

G. Cossu  
Pisa University & SNS


XXVI International Symposium on Lattice Field Theory  
Williamsburg, Virginia, USA, July 14-19 2008

In collaboration with:  
C. Bonati (Pisa), M. D'Elia (Genova),  
A. Di Giacomo (Pisa), C. Pica (BNL)

## **Outline**

- ✓ The order of the chiral transition in  $N_f=2$  QCD and the QCD phase diagram
- ✓ Prediction from effective models and previous literature
- ✓ Present evidence from lattice simulations
- ✓ (Extremely!) Preliminary new results
- ✓ Conclusion and discussion

# Phase diagram



## A test of first order scaling in $N_f=2$ QCD: progress report

The order of the transition at  $\mu_B=0$  is of great importance: the existence of a critical endpoint at  $T_E$  stems from the hypothesis of a crossover at the massless,  $\mu=0$  theory.

There is a general tendency to accept the crossover scenario in the real QCD case ( $N_f=2+1$  with physical quark masses): it has been shown (Y. Aoki, Z. Fodor, S. D. Katz and K. K. Szabo, *Phys. Lett. B* 643, 46 (2006); *Nature* 443, 675 (2006)) that the susceptibility of a possible order parameter for the transition (the chiral condensate) does not show any signal of growing with the spatial volume, till  $L_S \sim 6$  fm.

However is interesting to deeply address the problem because experimental evidences still lack.

Two flavor QCD in the light quark mass limit is an interesting case:

- there are theoretical predictions about the order (eff. Chiral models)
- no definite answer from lattice simulations yet

M. D'Elia, A. Di Giacomo, C. Pica, *PRD* 72, 114510 (2005)

G. C., M. D'Elia, A. Di Giacomo, C. Pica, arXiv:0706.4470

# Model predictions for chiral $N_f=2$

- $U_A(1)$  anomaly effective (no light  $\eta'$ ,  $c \neq 0$ ): effective model has a fixed point, i.e. **second order transition in the  $O(4)$  universality class or a first order.**
- $U_A(1)$  anomaly not effective (light  $\eta'$ ,  $c = 0$ ): no  $O(4)$  stable f. p.  
F. Basile, A. Pelissetto, E. Vicari, 2005  $\implies U(2)_L \otimes U(2)_R / U(2)_V$  or first order

Second order in the chiral limit  $\implies$  **crossover at small quark masses**

First order in the chiral limit  $\implies$  **first order in a small region around the chiral point**

# Determining the order: strategies

- Try the easiest (?) thing: look for **metastabilities and double peak structure** of the order parameter and of the energy density around the transition, i.e. coexistence of phases, which is a clear signature for first order. This search failed in the past leading to a preference for the second order scenario.
- Perform an accurate **Finite Size Scaling** analysis of various thermodynamical quantities around the chiral critical point to extract critical indexes ( $O(2)$  is expected rather than  $O(4)$  for the staggered fermion formulation).

	$y_h$	$y_t$	$\nu$	$\alpha$	$\gamma$
$O(4)$	<b>1.336(25)</b>	<b>2.487(3)</b>	<b>0.748(14)</b>	<b>-0.24(6)</b>	<b>1.479(94)</b>
$O(2)$	<b>1.496(20)</b>	<b>2.485(3)</b>	<b>0.668(9)</b>	<b>-0.005(7)</b>	<b>1.317(38)</b>
1st Order	<b>3</b>	<b>3</b>	<b>1/3</b>	<b>1</b>	<b>1</b>

# Finite Size Scaling

Approaching the transition the correlation length of the order parameter  $\xi$  goes large and one can write the following scaling ansatz:

**free energy density**  $\implies L/kT \simeq L_S^{-d} \phi(\tau L_S^{1/\nu}, am_q L_S^{y_h})$

**specific heat**  $\implies C_V - C_0 \simeq L_S^{\alpha/\nu} \phi_c(\tau L_S^{1/\nu}, am_q L_S^{y_h})$


**order parameter suscept.**  $\implies \chi \simeq L_S^{\gamma/\nu} \phi_\chi(\tau L_S^{1/\nu}, am_q L_S^{y_h})$

Technical difficulties:

- Simulations on large volumes and with light quark masses are necessary for a reliable f.s.s. analysis  $\implies$  huge computational power required
- f.s.s. behavior is given in terms of **two different scales** (two scaling variables).

No clear answer from previous literature (see our works for references).

# Previous work: second order check




$O(4)$  and  $O(2)$  are ruled out by our data. Notice that  $\alpha < 0$  for  $O(4)$



# Previous work: first order check

An approximate check for a first order scaling on the collected data was performed.





Indication that the transition **could be** first order. So we decided to give a chance to this hypothesis. But:

- where is the growth linear with the spatial volume expected for a first order transition at fixed mass?
- where are the double peaks?

# Further tests: 1<sup>st</sup> order direct check

G. C., M. D'Elia, A. Di Giacomo, C. Pica, arXiv:0706.4470



- Chiral susceptibility shows deviations, possibly due to the large mass range explored (up to 0.1) which could be outside the scaling region.
- Good scaling of the specific heat: not only the peak heights but also the widths are well described by the first order hypothesis.

# First order scaling analysis

Consider again the scaling law  $C_V - C_0 \simeq L_s^{\alpha/\nu} \phi_c(\tau L_s^{1/\nu}, \tau a m_q L_s^{y_h})$ .

- Continuous transition  $\implies L_s$  dependence must cancel as  $L_s \rightarrow \infty$  at finite  $m_q$ . The scaling function can be expanded in terms of  $1/(a m_q L_s^{y_h})$ : the leading term must be  $1/(a m_q L_s^{y_h})^{\alpha/\nu y_h} \implies$  **no discontinuity (no latent heat) at finite  $m_q$ .**
- First order chiral transition  $\implies$  a first order singularity is expected also at some  $m_q \neq 0$ , leading to a non-zero latent heat: **we can allow for a constant term** in the expansion in powers of  $1/(a m_q L_s^{y_h})$

$$C_V - C_0 \sim a m_q^{-1} \phi_c(\tau \vec{V}) + \mathbf{V} \phi'_c(\tau \vec{V})$$


In the second case the relative weight of the singular to the regular contribution is not known a priori, may be very small for small volumes and weak first order transitions.

## A test of first order scaling in $N_f=2$ QCD: progress report

There are various possibilities:


- There is really a first order transition which however is so weak that metastabilities will not show up but on very large, still unexplored volumes.
- We observe the “wrong” critical indexes because the scaling region around the chiral point is so small that the “correct”  $O(4)$  indexes will not show up but at very small, still unexplored quark masses.

In order to clarify the issue, we have judged worth dedicating a large numerical effort to a run at  $am_q=0.01335$  on a  $48^3 \times 4$  lattice (thanks to apeNEXT!)




That corresponds to  $m_\pi \sim$  twice the physical value and to a spatial size  $\sim 13-14$  fm.

# Preliminary results - Histories




We collected  $\sim 30k$  trajectories in total


# Preliminary results - Histories



# Preliminary results - Histories




# Preliminary results - Histories







# Preliminary results - Histograms




# Preliminary results - Histograms



# Preliminary results - Histograms



# Preliminary results - Histograms




Dated half June 2008.


# Preliminary results - Histograms

Spatial Plaquette Histogram

$m_q=0.01335$



# Preliminary results - Susceptibility



Shrinks with the correct factor but doesn't grow. We need more statistics. We hope to completely clarify this issue in the next months.

# Discussion and conclusions

**Conclusion 1:** With present UV cutoff effects ( $N_t=4$ , non-improved action) and within the present quark mass range a second order chiral transition in the  $O(4)$  (and  $O(2)$  and  $U(2)_L \otimes U(2)_R/U(2)_V$ ) seems to be excluded.

**Conclusion 2:** First order critical indexes seem to be preferred.  
**Preliminary:** we have some signals (**to be confirmed!!**) for a first order bistability at  $am_q=0.01335$ , however the bistability does not show up until  $L_s=12/T \sim 13-14$  fm. **LOW STATISTICS! Needs more investigation!**

Our results have been obtained with a quite large lattice spacing  $N_t=4$ , 0.3 fm (lattice spacing) and with a non-improved action. If our results will be confirmed on  $N_t=6$  and/or using an improved lattice action, then the crossover scenario must be changed.

 **Thank you!**

**Powered by:**

 **OpenOffice.org**

